


Managing Woody Debris



Leave it IN!

Woody debris should be left in a watercourse unless there is a very strong and well-supported case for its removal or repositioning. If reduced local water levels and flood duration is the primary rationale for LWD removal, the case should be supported by a hydraulic analysis.

Reposition

If a case for repositioning LWD is successfully made, a management proposal should be drawn up and circulated to all interested parties and relevant agencies for consultation. LWD needs to be pegged in at 20°-40°. This should improve the capacity of the channel to carry peak flows, while retaining a reasonable variety of low velocity habitats.

Reintroduction

The addition of LWD improves physical habitat and counteracts stream incision. 'Re-snagging' is now a common river rehabilitation technique in many parts of the world, in particular the USA, Australia and Canada. Introduced material should form the key pieces of stable debris accumulation:

- *Length: at least as long as the channel width*
- *Diameter: at least 0.1 metre or 5% of channel width (whichever is largest)*
- *LWD often needs to be securely keyed into the bed of the watercourse*



Possible Cases for Selective Removal

- *Debris accumulating on bridges and culverts, potentially resulting in localised flooding*
- *Navigation*
- *Urban Areas: Sewage litter and rubbish*
- *Confirmed long-term barrier to salmonids (BUT see page 14)*



Repositioned LWD

Catchment Strategies

Attitudes towards woody debris are changing in the UK. In many areas agencies are reducing their watercourse maintenance programmes. Interested parties are now in a position to develop strategies for the management of woody debris in river catchments. This can be carried out as a mapping exercise to identify both potential constraints and opportunities.



Riverine Woodland

Information gathering

- River Habitat Surveys (RHS), River Corridor Surveys and aerial photos provide data on the degree of bankside tree cover, LWD, debris dams, braided channels, islands and wet woodland.
- Local Environmental Records Centres maintain data on species, designated sites, nature reserves and County Wildlife Sites.
- UK and Local BAP partnerships have agreed targets for habitats like wet woodland and for flagship species such as *Lipsothrix* craneflies, white-dawed crayfish and Atlantic salmon.
- Agencies and local authorities have information on listed structures (bridges, weirs, water mills), bridges and culverts with poor capacity to handle peak flows, settlements in major flood zones, commercial forestry areas and community forests.

Promoting woody debris

- Do not disturb naturally occurring LWD unless it is vital for safety reasons or to protect investments (e.g. from flooding).
- Protect riparian corridors and floodplains to ensure that there will be a future source of woody debris. Influence agri-environment schemes in riparian areas and floodplains: promote the reduction, rotation or exclusion of livestock grazing, selective fencing of watercourses, buffer strips, planting schemes and natural regeneration areas. Avoid the uniform pollarding of willows and encourage a higher percentage to mature and collapse.
- When building, maintaining or improving roads and railways ensure that culverts and bridges have the capacity to allow flood flows and for LWD to pass through them.
- In the UK it will be many decades before the majority of watercourses once again begin to build up a resource of woody debris. Catchment strategies should help to identify suitable areas to reintroduce LWD in the short and medium term.



A benchmark lowland RHS site

Barrier or Boon for Fish?

Log jams that span a stream are often thought to create barriers to fish passage. As a result, many concerned woodland and fisheries managers have removed them. The irony is that, out of a desire to increase access for fish to habitat upstream, many efforts



to remove perceived 'blockages' are in fact detrimental to fish (and other wildlife) habitats downstream. Removing 'key pieces' of LWD can destabilise the remaining debris resulting in increased sedimentation of pools and gravels downstream. These pools may be reduced in size or lost altogether and gravels become unsuitable for fish spawning. Spawning gravels, previously held in place by stable LWD, can also be flushed downstream and dispersed.

Debris dams are generally observed and removed during normal flows. They are rarely observed during high flows when salmon and sea trout migration to spawning areas occurs. During high flows, debris dams can become submerged as water passes over the top or they can 'raft' upwards as water pushes from beneath. Fish are very adept at finding spaces above, below, around or within log jams. They have been doing it unaided for millennia!



Nevertheless, there are extreme instances where log jams become 'sealed' (perhaps by

forestry operations upstream or high silt burdens as a result of diffuse pollution) and prevent fish passage. Selective removal rather than complete removal is recommended. Ongoing monitoring of fish, other key species (eg otter and *Lipsothrix*), geomorphology (gravels, sediments and organic matter) and hydrology is important. Generally the removal of log jams to promote fish access needs to be carefully considered and weighed against the benefits summarised on pages 4-7.

Large Floodplain Rivers

LWD also has an important rôle to play in large floodplain rivers. Downed trees and living pieces of sprouting driftwood dramatically accelerate the process of island formation. If deposited 'living wood' (usually willows and poplars) is able to sprout and anchor to river sediments through the growth of adventitious roots, then the process of creating a pioneer island can begin. Sediments and gravel accumulate in the protected lee of the deposited LWD. These are good sites for willow regeneration and, in time, could help to produce a braided river channel.



River islands or braided channels are rare due to past land drainage works and a lack of naturally occurring LWD. Islands and their associated habitats support a range of biodiversity and it is time to identify areas where large rivers can be restored to their former character.



Case Studies

1. Black Water, New Forest National Park

Background The New Forest has the highest recorded density of debris dams in Britain. It contains some of the most important and rare wetlands in Europe including spectacular stands of riverine woodland, bog woodland and valley mires. However, many watercourses were re-sectioned and large areas drained for timber growing and livestock grazing.



A newly created log jam

Objectives To restore 600 hectares of wetland habitat, in part, by re-connecting watercourses with their former channels. To restore ten kilometres of damaged watercourse.



LWD introduced to the channel

Project Description Log jams have been introduced at strategic locations to help re-wet old stream courses.

Assessment The Black Water scheme is part of a completed Life3 Project. Monitoring has been carried out by PhD students. Results indicate that: (i) flood 'travel time' has been increased. Water now takes 35% longer to travel the length of the catchment due to storage on the restored wooded floodplains; (ii) there is an increase in channel diversity characterised by more woody debris and more pools.

Project Partners Forestry Commission, Environment Agency, English Nature, RSPB, Hampshire County Council, National Trust and the University of Southampton.

Cost Part of £2.9 million project

More info

www.newforestlife.org.uk/life3

www.geog.soton.ac.uk



One of the restored former channels

2. Tittesworth, Peak District National Park

Background Tittesworth Reservoir is owned and managed by Severn Trent Water. There are several fine examples of log jams and debris dams on the feeder streams that drain into the reservoir. However, one tributary in particular was identified as having been heavily modified in the past. This stream corridor became the focus of a habitat rehabilitation project.



Log weirs were installed at strategic locations and designed to create some ponded reaches upstream (but still allow fish passage). Raised water levels helped to re-wet approximately ten hectares of floodplain and a number of scrapes, ephemeral 'dragonfly' pools, 'ridge and furrow' reedbed shells and several more permanent ponds.



Park Authority, RSPB, Middlemarch Environmental, Landscape Matters.

Cost Approximately £25K for stream corridor work (part of £95K scheme)

More info n.mott@staffs-wildlife.org.uk

Objectives To increase habitat and species diversity at the site in line with Severn Trent Water's and the Peak Park's Biodiversity Action Plans. Use the scheme to inspire similar work at other sites as appropriate.

Description

700m of stream course were re-



Assessment

This imaginative and inspiring project has benefited a number of BAP target species. Monitoring has identified increases in breeding wildfowl and waders. Evidence of water voles was also detected for the first time.

Project Partners Severn Trent Water, Defra, Staffordshire Wildlife Trust, Environment Agency, Peak District National

3. River Trent at Wolsley Bridge, Staffordshire



Background Staffordshire Wildlife Trust (SWT) owns and manages the site. The River Trent had been engineered into a deep trapezoidal channel and effectively disconnected from its floodplain.

Objective To re-profile 340 metres of the River Trent along the inside of a large meander to help kickstart natural geomorphological processes including erosion and deposition and promote re-

alignment with its floodplain. To use the scheme as a demonstration of best practice to inspire similar work elsewhere.

Description The river re-profiling has been very successful. Shortly after completing the works a 'happy accident' involved the input of a mature beech tree which formed LWD in the restored channel. Liaison with the Environment Agency ensured that the LWD could be left in and monitored.



Assessment Monitoring of the LWD over a three year period has shown that (i) a short-term erosion pocket was created shortly after the beech tree entered the channel. The volume of the erosion pocket was approximately one and half times the volume of the submerged timber when the river was bank-full. It has since stabilised; (ii) No part of the LWD has moved more

than twenty metres downstream; (iii) The LWD has gradually accumulated smaller pieces of woody debris and has become surprisingly stable; (iv) common sandpiper, grey wagtail, kingfisher, homet and otter are among the species recorded using the LWD as a foraging, perching or spraint site.

Project Partners SWT, Environment Agency, Middlemarch Environmental

Cost Part of £21K project

More info n.mott@staffs-wildlife.org.uk



Reading

- Boyer, K.L., Berg, D.R. and Gregory, S.V. (2003) Riparian management for wood in rivers. pp. 407-420 In Gregory, S., Boyer, K. and A. Gurnell (Eds.) The Ecology and Management of Wood in World Rivers. American Fisheries Society Symposium 37
- Godfrey, A. (2003) A review of the invertebrate interest of coarse woody debris in England. Natural England Research Report Number **513** *
- Godfrey, A. (2000/2001/2005/2006) Surveys for the cranefly *Lipsothrix nigristigma* in 1999/2000/2003-2004/2005. Natural England Research Reports **351, 410** * / **Action for Invertebrates**
- Gurnell, A., Tockner, K., Edwards, P. and Petts, G (2005) Effects of deposited wood on biocomplexity of river corridors. Front Ecol Environ 2005; 3(7): 377-382 http://www.unc.edu/depts/geog/information/colloquia/Abstract_aGurnell.pdf
- Hughes, F.M.R. (ed.) (2003) The Flooded Forest: Guidance for policy makers and river managers in Europe on the restoration of floodplain forests. FLOBAR2, Department of Geography, University of Cambridge, UK
- Linstead, C. and Gurnell A.M. (1999) Large woody debris in British Headwater Rivers. Environment Agency Research & Development Technical Report **W 181** <http://publications.environment-agency.gov.uk/pdf/STR-W181-e-e.pdf> Summary Report: **W 185** <http://publications.environment-agency.gov.uk/pdf/STR-W185-e-e.pdf>
- Mackenzie, N.A. (1996) The Riparian Woodland Ecotone. Scottish Native Woods
- Opperman, J., Merenleder, A., and Lewis, D (2006) Maintaining Wood in Streams: A Vital Action for Fish Conservation. ANR University of California Publication 8157 <http://www.landstewards.org/otherDocs/MaintainingWoodInStreams.pdf>
- River Habitat Quality: the physical quality of streams and rivers in the UK and Isle of Man (1998). Environment Agency, Bristol
- Scottish Native Woods (2000) Restoring and Managing Riparian Woodlands

* These free publications can be ordered from Natural England's enquiry service enquiries@naturalengland.org.uk Tel. 0845 600 3078

Websites

Key Word Search: Large Woody Debris/Coarse Woody Debris/Large Wood/Snags/Re-snagging

- <http://www.sbes.stir.ac.uk/riversconference/>
- <http://panda.org/downloads/forests/deadwoodwithnotes.pdf>
- [http://www.forestresearch.gov.uk/pdf/FCGL002.pdf/\\$FILE/FCGL002.pdf](http://www.forestresearch.gov.uk/pdf/FCGL002.pdf/$FILE/FCGL002.pdf)
- <http://www.rivers.gov.au/acrobat/facts07.pdf>
- http://www.rivers.gov.au/acrobat/techguidelines/tech_guide_vol2_chapb.pdf
- http://www.dnr.state.oh.us/water/pubs/fs_st/stfs21.htm
- <http://www.ontariostreams.on.ca/OSRM/Tech6.pdf>
- <http://www.associationofrivertrusts.org.uk>

Credits

Author: Nick Mott; Design: Martin Adams

Photography: Nick Mott / Staffordshire Wildlife Trust

Additional Photography: p1 GCSlawson (debris dam), Sue Scott (Salmon), Darin Smith (Otter); p5 Sue Scott (Salmon); p6 Stuart Ball / JNCC (Mayfly), Steve Hewitt (Cranefly), David Iliff (Hoverfly); p8 Martin Cooper (R.Sow), Andrew Crawford, Environment Agency (R.Trent), Alan Jones, Environment Agency (R.Severn)

Map: Environment Agency / RHS

Publication data

Mott, N (2006) 'Managing Woody Debris in Rivers, Streams & Floodplains'. Staffordshire Wildlife Trust, UK

Printed by George Street Press, Stafford – Wildlife Trust Corporate Members

© Staffordshire Wildlife Trust and Natural England

01889-880100 www.staffs-wildlife.org.uk

Registered Charity No.259558



Water for Wildlife aims to take forward wetland conservation across the UK through partnership between The Wildlife Trusts, the Environment Agency, water companies, Water UK, and local communities. The Wildlife Trusts employ a network of Water for Wildlife staff throughout the country. To find out more about Water for Wildlife please visit www.waterforwildlife.org.uk.

The Wildlife Trusts is a partnership of 47 local Wildlife Trusts across the UK,. Our vision is 'an environment richer in wildlife for everyone' and we're the largest UK charity exclusively dedicated to conserving all our habitats and species, with a membership of more than 660,000 people including over 100,000 junior members. We campaign for the protection of wildlife and invest in the future by helping people of all ages to gain a greater appreciation and understanding of wildlife. Collectively, we also manage more than 2,500 nature reserves spanning over 80,000 hectares. To join, or for further information, phone The Wildlife Trusts on 0870 036 7711 or visit www.wildlifetrusts.org

The Environment Agency is the lead public body for protecting and improving the environment in England and Wales. The Agency works on the control and reduction of pollution of air, land and water, flood and coastal defence, water resource management, biodiversity, fisheries management, recreation and navigation. For more information visit www.environment-agency.gov.uk or for specific enquiries call 08708 506506.

***Please report all pollution incidents and fish kills immediately to
0800 80 70 60.***

Water UK The water industry is divided into twelve regional water and sewerage companies, with twelve smaller water-only companies. The companies are all members of Water UK, the organisation that represents the industry at national and European level. www.water.org.uk for more details.

Download PDF Version and Image Gallery from the Library section of www.staffs-wildlife.org.uk

Staffordshire Wildlife Trust wishes to acknowledge the financial support of Natural England in producing this booklet.

